



Published by the Society for Transparency, Openness, and Replication in Kinesiology under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, provided the original author and source are credited.

DOI:

10.51224/cik.2024.56

Subject Areas:


Physical Activity, Health, and Disease

Keywords:

Borderline personality disorder, physical activity, pattern, preferences, barriers, correlates

Editor:

Emily Budzynski-Seymour



Pattern, Preferences, Barriers, and Correlates of Self-Reported Physical Activity in Adults With Borderline Personality Disorder

An Online Survey in Western Countries

Samuel St-Amour^{1,2,*}, Lionel Cailhol^{2,3}, Josyane Lapointe^{1,2}, Déborah Ducasse^{4,5,6}, Gabrielle Landry³, and Paquito Bernard^{1,2}

¹Department of Physical Activity Sciences, Université du Québec à Montréal, Montréal, Québec, Canada

²Mental Health University Institute of Montreal Research Center, Montreal, Quebec, Canada

³Faculty of Medicine, Department of Psychiatry and Addictology, Université de Montréal, Montréal, Québec, Canada

⁴Department of Emergency Psychiatry and Post Acute Care CHU Montpellier, France

⁵IGF, Univ Montpellier, CNRS-INSERM, Montpellier, France

⁶Therapy Center for Mood and Emotional Disorders, Department of adult psychiatry, La Colombière, CHU Montpellier, France

*Correspondence: st-amour.samuel.2@courrier.uqam.ca

Borderline personality disorder (BPD) is characterized by an instability of self-image, interpersonal relationships, and emotions and is highly comorbid with other disorders. Physical activity has shown great results in treating these disorders. Physical activity intervention should be built considering the preferences and barriers of the targeted individuals. However, to this day no study analyzed the preferences and barriers to physical activity in individuals with BPD, which is the goal of this study. We used an online survey to question 192 adults with a self-reported diagnosis of BPD from Canada, France, the United States, England, Switzerland, and New Zealand. Participants complete 607 minutes of physical activity weekly on average. They prefer walking (66.7%), biking (33.3%), aquatic activities (29.0%), and running (24.2%). Their main barriers to physical activity are having a friend over, having other engagements, and recovering from an injury. Additionally, they prefer engaging in individually supervised physical activities outside, typically in long sessions of moderate intensity. Finally, a majority of participants are interested in receiving physical activity advice, but most did not receive any. The professionals from whom they would prefer to receive advice are trainers, psychiatrists, physical therapists, and psychologists. These findings are crucial for tailoring future physical activity interventions for adults with BPD.

Introduction

Borderline personality disorder (BPD) is the fourth most prevalent personality disorder in general population and the most prevalent in clinical settings (Gunderson et al., 2018). It is characterized by an instability of the self, individual goals, interpersonal relationships and affects (Gunderson et al., 2018). The biosocial development model of BPD suggests that emotion dysregulation is among the core components of the disorder and underlies some of its characteristic behaviours (Crowell et al., 2009). Moreover, emotion dysregulation has been associated with reduced quality of life, impaired daily functioning, and hindered therapeutic relationships (Gunderson et al., 2018). Individuals with BPD are also highly at risk to commit suicide with 83% having a history of suicide attempt (Soloff et al., 2002) and 8 to 10% dying from suicide (Biskin, 2015).

Those individuals also frequently present comorbid physical and mental disorders (El-Gabalawy et al., 2010; Shah & Zanarini, 2018). Mood disorders are the most frequent mental comorbid disorders in individuals with BPD with a lifetime prevalence of 96% (Shah & Zanarini, 2018). The main physical comorbid disorders in individuals with BPD include obesity, cardiovascular diseases and diabetes with one-year prevalence of 34%, 15%, and 9% respectively (Castle, 2019). Moreover, cardiovascular and metabolic disorders are among the greatest mortality causes with over 20% of death in this population (Cailhol et al., 2017; Kuo et al., 2018).

Among the treatments to address these different disorders, physical activity (PA) has been linked to improvements in many of their components and symptoms. The last Canadian treatment guidelines for mood disorders (the most prevalent comorbid disorder in individuals with BPD; Shah & Zanarini (2018)) included PA as mono- or adjunct therapy for every level of depression severity with the strongest level of proof (Ravindran et al., 2016). Finally, PA was also found efficient in preventing and reducing risk factors of obesity, cardiovascular diseases, and diabetes in individuals with mental disorder (Vancampfort et al., 2013). However, only one study to our knowledge analyzed the short-term effect of PA on BPD itself (St-Amour et al., 2022). No study examined the barriers, preferences and pattern of PA in adults with BPD as reported in a recent review (St-Amour et al., 2021; St-Amour et al., 2022).

To efficiently study PA's effects in individuals with BPD we can learn from the previous research done with individuals with other mental disorders. Among the many challenges of studying PA effects in individuals with mental disorders, dropout and adherence are among the greatest (Stubbs & Rosenbaum, 2018). Indeed, dropout rates in studies analyzing PA's effect in individuals with mental disorders may be as high as 90% and are heavily influenced by study and intervention characteristics (Firth et al., 2015). Moreover, participants who completed those studies did not necessarily attend all sessions. Attendance widely varied in these studies ranging from 30% to 100% of the planned seances (Firth et al., 2015). It is therefore important to provide optimal conditions to reach and more importantly keep participants in future PA studies. The latest guidelines for PA interventions in individuals with mental disorders suggest tailoring those interventions in light of the preferences and barriers of the targeted individuals or population (Romain & Bernard, 2018). It is therefore important to identify those preferences and barriers in individuals with BPD, which has not yet been done to our knowledge, to plan effective interventions. Since BPD is a more relational disorder than other mental disorders in its nature (Gunderson et al., 2018), preferences regarding PA is done and main barriers preventing PA participation could differ from populations with other mental disorders.

There is a growing interest in research for PA in individuals with BPD. However, to our knowledge there are very few observational studies (St-Amour et al., 2022) and only one experimental study (St-Amour et al., 2022) addressing this topic. To develop well-adapted PA interventions for adults with BPD in the future, we need to identify the most important factors associated with adherence.

Being the first study in this population, the aims are to: 1- describe the self-reported PA level of adults with BPD; 2- describe the main preferences regarding type, location, intensity, supervision, and advice related to PA in adults with BPD; 3-describe the main barriers to practise PA in adults with BPD; and 4-examine the main sociodemographic and health variables associated to the self-reported level of PA in adults with BPD.

Methods

This online cross-sectional study in western countries was performed using the LimeSurvey platform hosted on the Université du Québec à Montréal's servers. The survey was promoted online in Canada, France, England, the United States, New Zealand, Switzerland, Belgium, and Australia, where the research team had research collaborators, with YouTube videos, a Facebook page and posts on Facebook groups, forums, and chat groups dedicated to adults with BPD (with the permission of the respective administrators). Psychiatrists in France and Canada helped promote this survey by sharing it with their patients, their colleagues, and in different networks regrouping patients and professionals working with patients with BPD. To be included, participants had to report: 1-being at least 18 years old, 2-living in Canada, France, England, the United States, New Zealand, Switzerland, Belgium, or Australia, and 3-having received a BPD diagnosis from a healthcare professional. All participants had to read and agree to the online information and consent form and were given the opportunity to contact the research team prior to filling-out the survey. They were warned that the survey may address sensitive topics, and local help resources were given to participants according to their reported country of residence at the bottom of each page (Batterham, 2014; Choi et al., 2017). This study has been approved by the ethics boards from the Eastern Montreal Integrated University Health and Social Services Centres (2021-2330) and the Université du Québec à Montréal (3997_e_2021).

Questionnaires

Every questionnaire used in this survey has been previously validated in English and in French. All items and questionnaires in English and French are available in open access (<https://osf.io/5u6am/>). Questionnaires included in this survey relate to PA habits and the main correlates of PA found in the literature (income, education level, disorder severity and duration, medication use, substance use, and suicide ideations; St-Amour et al. (2021), Vancampfort et al. (2011), Vancampfort et al. (2018)).

Sociodemographic Characteristics

The following sociodemographic characteristics have been collected: country of residence, sex at birth, age, education level, marital status, height, weight, and household income. Participants have also been asked about their psychiatric follow-up duration, their psychotropic medication use, their comorbid disorders and the numbers of past mental disorder hospitalization and suicide attempts. Afterward, they had to rate their social status with the MacArthur Scale (Adler et al., 2008).

Clinical Characteristics

Participants then filled out the Borderline Symptoms List-Short Form (BSL-23) validated to measure the presence and severity of symptoms attributed to BPD (Bohus et al., 2008). Then, they filled the Beck Depression Inventory-Short Form (BDI-SF) validated to measure the clinical depression risk in adults with BPD. A score of 10 and higher indicate the presence of depression (Furlanetto et al., 2005). Difficulties in emotion regulation have been assessed with the Difficulties in Emotion Regulation Scale-Short Form (DERS-18) that have been validated in adults with BPD (Victor & Klonsky, 2016).

Health Behaviours

Substance use disorders have been assessed with the Alcohol, Smoking and Substance Involvement Screening Test (ASSIST). This questionnaire developed and validated by the World Health Organization (WHO) measures substance use disorders related to tobacco, alcohol, cannabis, cocaine, amphetamines, inhalants, sedatives, hallucinogens, and opioids. A score is obtained for each substance by adding the score from each question. A score of 4 and more (11 and more for alcohol) indicate a moderated substance use disorder and a score of 27 and more (including alcohol) indicate a severe substance use disorder (WHO ASSIST Working Group, 2002). Participants' insomnia was assessed with the Insomnia Severity Index (ISI) in which a score higher than 15 indicating a risk of clinical insomnia (Bastien, 2001). PA was assessed using the Global Physical Activity Questionnaire (GPAQ) developed by the WHO to measure work, leisure, travel, and total PA (Armstrong & Bull, 2006). The perceived built environment was rated using the Instrument for Assessing Levels of Physical Activity and Fitness (ALPHA) environmental questionnaire (Spittaels et al., 2009).

Physical Activity Preferences and Barriers

Preferences regarding type, intensity, context, supervision, and advice of PA and main barriers to PA have been assessed using questionnaires from previous studies in individuals with mental disorders (Abrantes et al., 2011; Romain, Longpré-Poirier, et al., 2020). For preference regarding the type of PA, participants had to select up to 5 specific activity among a list of 24 different choices. For preference regarding the person from whom they wish to receive PA advice, participants had to rank their 3 most preferred persons. For the full list of suggested activity, see supplementary material (Table S1). For every other question regarding preference, participant had to choose the most appropriate answer to their situation.

Statistical Analysis

Descriptive statistics (N, %, mean) are used to describe the sociodemographic characteristics, the substance use disorder, insomnia, and PA level, preferences, and barriers in this sample. A Poisson multivariate regression was done to identify the main sociodemographic and clinical factors associated with weekly PA level. A graph presenting the distribution of PA level among participants is available in supplementary material (Figure S1). The age, social level, education level, household income, body mass index, emotion regulation difficulties, BPD symptoms, tobacco and alcohol use disorders, depression, past suicide attempts, psychotropic medication use and self-efficacy were included in the model. These independent variables were selected based on previous papers examining the determinants of PA in adults with mental disorders (Schuch et al., 2017; Vancampfort et al., 2011; Vancampfort et al., 2015). The R function stepAIC was used with an initial full model and combined with a forward and backward selection (Zhang, 2016). The final model has the lowest Akaike information criterion.

The multicollinearity of this model has been tested by calculating the variance inflation factor (VIF). No predictors had VIF values greater than 10 after the stepwise variable selection procedure. Statistical analyses were done with the R software version 4.2 with the libraries “ggstaplot”, “stargazer”, “MASS”, and “summarytools” (Patil, 2021; Zhang, 2016). Research materials, data and R codes are available in open access on Open Science Framework (<https://osf.io/c3gvx/>).

Results

The online survey was completed by 288 participants who reported having a BPD diagnosis, but only 192 filled out the survey at least up to the GPAQ providing PA data. Therefore, the results are based on this sample of 192 that completed the GPAQ. Of those 192 participants, 96 were living in Canada, 78 were living in France, 9 in the United States, 7 in England and 2 in Switzerland. Description of the sample is presented in Table 1.

Table 1. Descriptive data of subjects

Variables	N (%)
Sex at birth	
Women	164 (85.9)
Age	
<35 years	111 (57.8)
Education	
< University education	118 (61.5)
Social Status	
Subjective Social Status M (SD)	4.5 (2.0)
Country of residence	
Canada	96 (50.0)

Variables	N (%)	
	France	78 (40.6)
	United States	9 (4.7)
	England	7 (3.6)
	Switzerland	2 (1.0)
Income		
	<20,000	71 (37.0)
	20,000-39,999	36 (18.8)
	40,000-59,999	21 (10.9)
	60,000-79,999	24 (12.5)
	80,000-99,999	10 (5.2)
	100,000	8 (4.2)
	Don't know	12 (6.2)
	No response	10 (5.2)
Marital status		
	Single	95 (47.4)
	Civil union	49 (25.5)
	Married	23 (12.0)
	Divorced	19 (9.9)
	Widowed	2 (1.0)
	Other	4 (2.1)
Stature		
	Body mass index M (SD)	26.4 (8.1)
Health Status		
	Self-reported diagnosis of physical disease	42 (21.9)
	Self-reported diagnosis of mental disorder	101 (52.6)
	Psychotropic medication	140 (72.9)
	Currently in psychotherapy	95 (47.4)
	Have been hospitalized	113 (58.9)
	Attempted suicide	126 (65.6)
Total substance involvement		
	High	111 (57.8)
	Moderate	20 (10.4)
	None	61 (31.8)
Psychological Measures		
	Insomnia severity index M (SD)	13.2 (6.5)

Variables	N (%)
Borderline symptoms list-short form M (SD)	1.9 (0.9)
Beck Depression inventory	16.0 (7.7)
Difficulty in emotion regulation scale M (SD)	59.9(13.5)

Note: continuous variables represented as mean (M) and standard deviation (SD)

Levels of Physical Activity

In this sample, 122 (65.9%) participants declared being physically active according to the guidelines from the WHO (completing at least 150 minutes of PA weekly; WHO. Regional Office for Europe & United Nations Economic Commission for Europe, 2022). On average, participants complete 165.5 (SD = 282.1) minutes of travel-related PA, 315.2 (SD = 685.7) minutes of work-related PA, and 126.0 (SD = 224.9) minutes of leisure-related PA for a total of 606.7 (SD = 766.2) minutes of PA weekly including all three domains. In this sample, men performed significantly more PA than women ($p = 0.002$). For detailed results comparing total PA levels according to body mass index, sex, level of education, age, and country, see supplementary material (Figure S2).

Barriers

Barriers preventing participants from engaging in PA were measured with self-efficacy scales from 0 to 100. For each barrier, participants had to rate their self-efficacy to engage in PA when encountering said barrier. Scores were reversed for clarity, therefore a higher score means a more important barrier. The PA barriers are illustrated in Figure 1. For detailed barriers according to age, education level, sex, body mass index, and country, see supplementary material (Figures S3-S7).

Note: The higher the mean indicator (vertical black line) on the abscissa, the greater the barrier. The different curves represent the distribution of data for each barrier. The barriers in order from the top are: “Being tired”, “Feeling pressure at work”, “Recovering from an injury”, “Being in a bad mood”, “Having personal problems”, “Having more interesting things to do”, “Without the support of friends or family”, “Having other engagement”, “Feeling unwell”, “Having friends at home”, “Feeling anxious”, “Not reaching previously fixed training goals”, “Having family problems”, “During vacations”.

Preferences

The detailed frequency to which each PA was declared as being among one’s preferred are represented in Figure 2. For detailed PA preferences according to body mass index, sex, country, education level, and age, see supplementary material (Figures S8-S12).

The details for preferred modalities and contexts of PA are presented in Figure 3.




Figure 1: Self-efficacy to do PA when encountering barriers.




Figure 2: Most frequently preferred physical activity.







Figure 3: Preferred modalities and context of physical activity. Note: BPD = borderline personality disorder; The sum of the % may not be equal to 100 due to missing data not included.

Advice


The details of the preferences regarding PA advice are presented in Figure 4.

Note: For the PA advisor, the scores for each professional were calculated by asking participants to indicate which 3 professionals they prefer receiving advice from with the numbers 1 to 3 (1 = 3rd choice; 2 = 2nd choice; 3 = 1st choice). All professional without number were attributed the number 0. The mean score for each professional (including 0s) were computed, reversed for clarity (making a mean score closer to 1 a higher preference for that professional) and reported here.

Received advices about physical activity



Mode of delivery



Autonomous of physical activity session



Self-efficacy level for participation to physical activity for BPD




Figure 4: Preference regarding physical activity advice.

Factors Associated with Physical Activity

Seven univariate outliers have been excluded because they had an exceptional self-reported PA level (i.e., more than 3500 min per week). A bivariate regression correlation table is presented in Figure S13 as supplementary material to show existing correlation in included variables. In this sample, the total level of PA is associated to the age, the social level, the education level, the income, the body mass index, the level of BPD symptoms, having a tobacco use disorder, the level of depression, the number

of psychotropic medications used and the self-efficacy to PA. The total score of DERS-18 and ALPHA questionnaire were initially in the full model, but they have been excluded during the variables selection presented in statistical section. For all the details see Table 2.

Table 2. Factors associated to level of physical activity

	Factors associated with total self-reported PA (SD)
Age	-0.005 (0.0003)*
Perceived social status	0.06 (0.002)*
University education	-0.23 (0.007)*
Income >40,000	-0.31 (0.007)*
BMI	0.006 (0.0004)*
BPD severity	0.27 (0.006)*
Tobacco SUD	0.38 (0.007)*
Alcohol SUD	0.03 (0.007)*
Depression	-0.04 (0.001)*
Psychotropic medication	0.09 (0.003)*
Self-efficacy	0.01 (0.002)*
Observations	180
Adjusted R ²	0.12
Log Likelihood	-66656
Akaike information criterion	133338

Note: BMI = body mass index;

BPD = borderline personality disorder;

SUD = substance use disorder.

*p < 0.01

Discussion

This is the first study to describe the PA level, preferences, barriers and correlates in a sample of adults with BPD. Although online studies pose many challenges, they are efficient and doable in adults with BPD (DeShong & Tucker, 2019; Lawn & McMahon, 2015).

Level of Physical Activity

First, we observe a relatively high level of PA in this sample. With a mean of more than 600 minutes weekly, participants are on average more active than the general population (about 340 minutes weekly; Colley et al. (2018)) and individuals with mental disorders (about 270 minutes weekly; Vancampfort et al. (2017)). With a little over 60%, this sample presents a similar proportion of active individuals as the general population (about 60%; Colley et al. (2018)) but a bigger proportion than individuals with other mental disorders (about 45%; Vancampfort et al. (2017)). These data are surprising and might indicate that the mean PA level is skewed by extreme individuals with large PA volume. However, the present data are difficult to put in perspective because of the absence of studies reporting PA level in this population (St-Amour et al., 2022).

Barriers

The main barriers to PA reported in this study, as measured by the lowest reported self-efficacy to overcome them, are having friends over, having other engagements, recovering from an injury, having something else more interesting to do and feeling too tired. In comparison, individuals with severe mental disorders or substance use disorder reported lack of motivation, fatigue, having no one to engage in PA with, not having enough energy, would not be able to keep up, and lack of financial resources as barriers to PA (Abrantes et al., 2011; Romain, Longpré-Poirier, et al., 2020). The barriers reported here were then somewhat different from those reported in other populations with mental disorders with the exception of lack of energy/feeling tired. Surprisingly, lacking support or not having anyone to engage in PA with was the least important barrier in adults with BPD but was among the main barriers in populations with other mental disorders (Abrantes et al., 2011; Romain, Longpré-Poirier, et al., 2020). This difference might be explained by the relational nature of the personality disorder. Indeed, one of the main characteristics of BPD is the difficulty in maintaining interpersonal relationships (Euler et al., 2021). These difficulties might lead individuals with BPD to prefer engaging in PA alone and therefore not considering lacking support or not having someone to do PA with as a major barrier to PA. Moreover, individuals with BPD tend to be more impulsive and to prefer quick smaller rewards than delayed larger rewards than healthy individuals and those with other mental disorders (Gunderson et al., 2018). Therefore, they might be tempted to engage in quickly rewarding activities rather than PA. However, more studies are needed to thoroughly understand this difference.

Preferences

The most frequently reported favourite PAs are somewhat consistent with the findings from populations with other mental disorders (Abrantes et al., 2011; Romain, Longpré-Poirier, et al., 2020). Indeed, walking is the preferred PA regardless of the studied population (Abrantes et al., 2011; Romain, Longpré-Poirier, et al., 2020). The clear preference of walking as a PA might be attributed to it being practical, self-paced and controlled, inexpensive, and not needing a lot of resources (Abrantes et al., 2011).

As mentioned before, participants in this study clearly prefer engaging in PA alone than with other people. This result differs greatly from what is seen in other populations with mental disorder having mostly no preference for doing PA alone or in groups. This difference might also be explained by the interpersonal difficulties experienced by individuals with BPD (Euler et al., 2021).

In this sample, most participants prefer supervised PA compared to unsupervised PA. This result is also surprising and in opposition to the findings observed in other populations. Indeed, in populations with other mental disorders, there is either a clear preference for unsupervised PA or no clear preference regarding supervision (Abrantes et al., 2011; Romain, Longpré-Poirier, et al., 2020). This result is unexpected considering previously observed results and interpersonal difficulties of individuals with BPD. However, this difference might be explained by the perception of the different relationships in play. Individuals with BPD tend to feel closer from those peripheral to their social network and farther from the more central individuals (Beeney et al., 2018). Therefore, they might feel closer to an outside individual like a kinesiologist or physical therapist supervising their PA session but would not want to share these moments with friends or relatives.

Advice

An overwhelming majority of participants declared being interested or maybe interested in receiving PA advice. However, a majority also declared not having received PA advice. Since all participants reportedly received BPD diagnosis from a healthcare professional, this lack of advice regarding PA is alarming considering two evidence-based BPD treatments suggest PA in their official guidelines (Blum et al., 2008; Linehan, 2014). Moreover, participants declared trusting healthcare professionals in giving them advice about PA, but preferred trainers, psychiatrists and physical therapists. It would therefore be important to include kinesiologists and PA professionals in multidisciplinary teams taking care of individuals with BPD. Future research may look at different PA promotion interventions in this population specifically (motivational interviews, messaging, group intervention, etc.).

Correlates of Physical Activity

In this study, sociodemographic and clinical variables have some opposite correlation to PA level with what is observed in previous systematic reviews (Vancampfort et al., 2011). Indeed, age, education level, and household income are negatively correlated and social status is positively correlated to PA level. Also, body mass index, BPD symptoms, tobacco use disorder, alcohol use disorder and number of psychotropic medications are positively correlated. Higher levels of depression is associated with lower PA level. Finally, self-efficacy level is also positively linked to PA level which is also observed in a recent meta-analysis (Cabassa et al., 2020). Among these correlations, the most surprising are those with the education level, the household income, the level of BPD symptoms, the tobacco and alcohol use disorder, and the number of psychotropic medications. The association between lower income and education level, and higher PA level in the present sample might be explained by a greater proportion of work-related PA in individuals with lower income and education level (Statistics Canada et al., 2020). Moreover, work-related PA is more strongly correlated with the total PA volume than leisure time PA and most studies reporting PA level only report leisure time PA (Vancampfort et al., 2011). Health-related surprising correlates (body mass index, BPD symptoms, substance use disorders and medication) might be explained by the nature of psychiatric comorbid disorders in our participants (St-Amour et al., 2021; Vancampfort et al., 2011). Following the suggestion of a reviewer, a sensitivity analysis was realized using only leisure time PA as dependent variable, confirming the supposition regarding education level and household income's surprising relation to PA. Moreover, relations between BMI and alcohol use disorder, and leisure time PA have also been reversed which is more in line with results from the literature. Finally, past suicide attempts became significantly associated with leisure time PA. For full details of these sensitivity analysis, see supplemental material (Table S2).

Limitations

However, this study suffers from some limitations. First, BPD diagnosis was self-reported making it hard to ensure this sample is composed solely of adults with BPD. This sample is also small considering the observational design adopted. Second, this study could have attracted more active than inactive participants due to its theme (i.e., PA) subjecting it to recruitment bias and increasing PA levels artificially. Using self reported PA measures may subject our results to recall bias and therefore affect the validity of the data. Future studies could use representative national surveys reporting both BPD diagnosis and objectively measured PA. However, to our knowledge, no such survey is available at the moment. Moreover, the unequal recruitment between countries biases the results with an over-representation of Canada and France limiting the generalizability in other countries. The online nature of this survey also poses some limitations. Indeed, online surveys are only accessible to those with access to a computer and sufficient informatics literacy to complete them. In addition, the cross-sectional design of this study prevents us from establishing causality relations between PA and its correlates. Finally, participants may have developed questionnaire fatigue because of the length of the survey, but this was not measured nor analyzed.

Conclusions

To our knowledge, this is the first study analyzing PA level, preferences, barriers, and correlates in adults with BPD. This information is primordial in developing future studies analyzing PA's medium- to long-term effect in adults with BPD. With treatment guidelines already suggesting the use of PA (Blum et al., 2008; Linehan, 2014), and indirect evidence indicating potential benefits of PA in alleviating BPD symptoms (Mehren et al., 2020), there is an urgent need for data analyzing the effect of PA on the different components of BPD to test this hypothesis. Pending future results, PA could be used by healthcare professionals (mostly kinesiologists, psychiatrists, and psychologists) to treat or alleviate comorbid disorders in this population. In doing so, they should base their intervention on the present barriers and preferences (i.e., walking, individual intervention, supervised sessions, outdoor, etc.) to ensure greater adherence (Romain, Bernard, et al., 2020). Future studies should also use nationally representative surveys to answer these questions and aim to compare these results with those from individuals with other mental disorders.

Additional Information

Data Accessibility

Data is available via the Open Science Framework account (<https://doi.org/10.17605/OSF.IO/C3GVX/>)
DOI: 10.17605/OSF.IO/C3GVX

Author Contributions

- Contributed to conception and design: SS, LC, PB
- Contributed to acquisition of data: SS, LC, DD, GL
- Contributed to analysis and interpretation of data: SS, PB
- Drafted and/or revised the article: SS, LC, JL, DD, GL, PB
- Approved the submitted version for publication: SS, LC, JL, DD, GL, PB

Conflict of Interest

The authors have no conflict of interest to declare.

Funding

This study was supported by PB's Junior-1 research grant from the *Fonds de Recherche du Québec - Santé*.

Acknowledgments

We would like to thank every participant for their valuable contribution in realizing this study and everyone who got involved in the promotion of the survey and the recruitment process.

Preprint

The pre-publication version of this manuscript can be found on MedRxiv at <https://doi.org/10.1101/2022.05.24.22275513> (DOI: 10.1101/2022.05.24.22275513)

References

- Abrantes, A. M., Battle, C. L., Strong, D. R., Ing, E., Dubreuil, M. E., Gordon, A., & Brown, R. A. (2011). Exercise preferences of patients in substance abuse treatment. *Mental Health and Physical Activity, 4*(2), 79–87. <https://doi.org/10.1016/j.mhpa.2011.08.002>
- Adler, N., Singh-Manoux, A., Schwartz, J., Stewart, J., Matthews, K., & Marmot, M. G. (2008). Social status and health: A comparison of British civil servants in Whitehall-II with European- and African-Americans in CARDIA. *Social Science and Medicine, 66*(5), 1034–1045. <https://doi.org/10.1016/j.socscimed.2007.11.031>
- Armstrong, T., & Bull, F. (2006). Development of the World Health Organization Global Physical Activity Questionnaire (GPAQ). *Journal of Public Health, 14*(2), 66–70. <https://doi.org/10.1007/s10389-006-0024-x>
- Bastien, C. (2001). Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Medicine, 2*(4), 297–307. [https://doi.org/10.1016/s1389-9457\(00\)00065-4](https://doi.org/10.1016/s1389-9457(00)00065-4)
- Batterham, P. J. (2014). Recruitment of mental health survey participants using Internet advertising: content, characteristics and cost effectiveness. *International Journal of Methods in Psychiatric Research, 23*(2), 184–191. <https://doi.org/10.1002/mpr.1421>
- Beeney, J. E., Hallquist, M. N., Clifton, A. D., Lazarus, S. A., & Pilkonis, P. A. (2018). Social disadvantage and borderline personality disorder: A study of social networks. *Personality Disorders: Theory, Research, and Treatment, 9*(1), 62–72. <https://doi.org/10.1037/per0000234>
- Biskin, R. S. (2015). The Lifetime Course of Borderline Personality Disorder. *The Canadian Journal of Psychiatry, 60*(7), 303–308. <https://doi.org/10.1177/070674371506000702>
- Blum, N., M.S.W., St. John, D., M.A., P.A., Pfohl, B., M.D., Stuart, S., M.D., McCormick, B., M.A., Allen, J., Ph.D., Arndt, S., Ph.D., & Black, D. W., M.D. (2008). Systems Training for Emotional Predictability and Problem Solving (STEPPS) for Outpatients With Borderline Personality Disorder: A Randomized Controlled Trial and 1-Year Follow-Up. *American Journal of Psychiatry, 165*(4), 468–478. <https://doi.org/10.1176/appi.ajp.2007.07071079>
- Bohus, M., Kleindienst, N., Limberger, M. F., Stieglitz, R.-D., Domsalla, M., Chapman, A. L., Steil, R., Philipsen, A., & Wolf, M. (2008). The Short Version of the Borderline Symptom List (BSL-23): Development and Initial Data on Psychometric Properties. *Psychopathology, 42*(1), 32–39. <https://doi.org/10.1159/000173701>
- Cabassa, L. J., Stefancic, A., Wang, X., Guo, S., Lu, N. Y., & Weatherly, C. (2020). Correlates of Physical Activity and Cardiorespiratory Fitness in Racially and Ethnically Diverse People with Serious Mental Illness in Supportive Housing. *Community Mental Health Journal, 56*(6), 1139–1152. <https://doi.org/10.1007/s10597-020-00610-x>
- Cailhol, L., Pelletier, É., Rochette, L., Laporte, L., David, P., Villeneuve, É., Paris, J., & Lesage, A. (2017). Prevalence, Mortality, and Health Care Use among Patients with Cluster B Personality Disorders Clinically Diagnosed in Quebec: A Provincial Cohort Study, 2001-2012. *The Canadian Journal of Psychiatry, 62*(5), 336–342. <https://doi.org/10.1177/0706743717700818>
- Castle, D. J. (2019). The complexities of the borderline patient: how much more complex when considering physical health? *Australasian Psychiatry, 27*(6), 552–555. <https://doi.org/10.1177/1039856219848833>
- Choi, I., Milne, D. N., Glozier, N., Peters, D., Harvey, S. B., & Calvo, R. A. (2017). Using different Facebook advertisements to recruit men for an online mental health study: Engagement and selection bias. *Internet Interventions, 8*, 27–34. <https://doi.org/10.1016/j.invent.2017.02.002>
- Colley, R. C., Butler, G., Garriguet, D., Prince, S. A., & Roberts, K. C. (2018). Comparison of self-reported and accelerometer-measured physical activity in Canadian adults. *Health Reports, 29*(1), 3–15. <https://pubmed.ncbi.nlm.nih.gov/30566204/>
- Crowell, S. E., Beauchaine, T. P., & Linehan, M. M. (2009). A biosocial developmental model of borderline personality: Elaborating and extending linehan's theory. *Psychological Bulletin, 135*(3), 495–510. <https://doi.org/10.1037/a0015616>
- DeShong, H. L., & Tucker, R. P. (2019). Borderline personality disorder traits and suicide risk: The mediating role of insomnia and nightmares. *Journal of Affective Disorders, 244*, 85–91. <https://doi.org/10.1016/j.jad.2018.10.097>
- El-Gabalawy, R., Katz, L. Y., & Sareen, J. (2010). Comorbidity and Associated Severity of Borderline Personality Disorder and Physical Health Conditions in a Nationally Representative Sample. *Psychosomatic Medicine, 72*(7), 641–647. <https://doi.org/10.1097/psy.0b013e3181e10c7b>

- Euler, S., Nolte, T., Constantinou, M., Griem, J., Montague, P. R., Fonagy, P. (2021). Interpersonal Problems in Borderline Personality Disorder: Associations With Mentalizing, Emotion Regulation, and Impulsiveness. *Journal of Personality Disorders*, *35*(2), 177–193. https://doi.org/10.1521/pedi_2019_33_427
- Firth, J., Cotter, J., Elliott, R., French, P., & Yung, A. R. (2015). A systematic review and meta-analysis of exercise interventions in schizophrenia patients. *Psychological Medicine*, *45*(7), 1343–1361. <https://doi.org/10.1017/s0033291714003110>
- Furlanetto, L. M., Mendlowicz, M. V., & Romildo Bueno, J. (2005). The validity of the Beck Depression Inventory-Short Form as a screening and diagnostic instrument for moderate and severe depression in medical inpatients. *Journal of Affective Disorders*, *86*(1), 87–91. <https://doi.org/10.1016/j.jad.2004.12.011>
- Gunderson, J. G., Herpertz, S. C., Skodol, A. E., Torgersen, S., & Zanarini, M. C. (2018). Borderline personality disorder. *Nature Reviews Disease Primers*, *4*(1). <https://doi.org/10.1038/nrdp.2018.29>
- Kuo, C.-J., Chen, W.-Y., Tsai, S.-Y., Chen, P.-H., Ko, K.-T., & Chen, C.-C. (2018). Excessive mortality and causes of death among patients with personality disorder with comorbid psychiatric disorders. *Social Psychiatry and Psychiatric Epidemiology*, *54*(1), 121–130. <https://doi.org/10.1007/s00127-018-1587-x>
- Lawn, S., & McMahon, J. (2015). Experiences of care by Australians with a diagnosis of borderline personality disorder. *Journal of Psychiatric and Mental Health Nursing*, *22*(7), 510–521. <https://doi.org/10.1111/jpm.12226>
- Linehan, M. (2014). In *DBT? Skills Training Manual*. Guilford Publications.
- Mehren, A., Reichert, M., Coghill, D., Müller, H. H. O., Braun, N., & Philipsen, A. (2020). Physical exercise in attention deficit hyperactivity disorder – evidence and implications for the treatment of borderline personality disorder. *Borderline Personality Disorder and Emotion Dysregulation*, *7*(1). <https://doi.org/10.1186/s40479-019-0115-2>
- Patil, I. (2021). Visualizations with statistical details: The “ggstatsplot” approach. *Journal of Open Source Software*, *6*(61), 3167. <https://doi.org/10.21105/joss.03167>
- Ravindran, A. V., Balneaves, L. G., Faulkner, G., Ortiz, A., McIntosh, D., Morehouse, R. L., Ravindran, L., Yatham, L. N., Kennedy, S. H., Lam, R. W., MacQueen, G. M., Milev, R. V., Parikh, S. V. (2016). Canadian Network for Mood and Anxiety Treatments (CANMAT) 2016 Clinical Guidelines for the Management of Adults with Major Depressive Disorder. *The Canadian Journal of Psychiatry*, *61*(9), 576–587. <https://doi.org/10.1177/0706743716660290>
- Romain, A. J., & Bernard, P. (2018). Behavioral and Psychological Approaches in Exercise-Based Interventions in Severe Mental Illness. In *Exercise-Based Interventions for Mental Illness: Physical Activity as Part of Clinical Treatment* (pp. 187–217). Academic Press. <https://doi.org/10.1016/B978-0-12-812605-9.00010-1>
- Romain, A. J., Bernard, P., Akross, Z., St-Amour, S., Lachance, J.-P., Hains-Monfette, G., Atoui, S., Kingsbury, C., Dubois, E., Karelis, A. D., & Abdel-Baki, A. (2020). Motivational theory-based interventions on health of people with several mental illness: A systematic review and meta-analysis. *Schizophrenia Research*, *222*, 31–41. <https://doi.org/10.1016/j.schres.2020.05.049>
- Romain, A. J., Longpré-Poirier, C., Tannous, M., & Abdel-Baki, A. (2020). Physical activity for patients with severe mental illness: Preferences, barriers and perceptions of counselling. *Science and Sports*, *35*(5), 289–299. <https://doi.org/10.1016/j.scispo.2020.03.005>
- Schuch, F., Vancampfort, D., Firth, J., Rosenbaum, S., Ward, P., Reichert, T., Bagatini, N. C., Bgeginski, R., & Stubbs, B. (2017). Physical activity and sedentary behavior in people with major depressive disorder: A systematic review and meta-analysis. *Journal of Affective Disorders*, *210*, 139–150. <https://doi.org/10.1016/j.jad.2016.10.050>
- Shah, R., & Zanarini, M. C. (2018). Comorbidity of Borderline Personality Disorder. *Psychiatric Clinics of North America*, *41*(4), 583–593. <https://doi.org/10.1016/j.psc.2018.07.009>
- Soloff, P. H., Lynch, K. G., & Kelly, T. M. (2002). Childhood Abuse as a Risk Factor For Suicidal Behavior in Borderline Personality Disorder. *Journal of Personality Disorders*, *16*(3), 201–214. <https://doi.org/10.1521/pedi.16.3.201.22542>
- Spittaels, H., Foster, C., Oppert, J.-M., Rutter, H., Oja, P., Sjöström, M., & De Bourdeaudhuij, I. (2009). Assessment of environmental correlates of physical activity: development of a European questionnaire. *International Journal of Behavioral Nutrition and Physical Activity*, *6*(1). <https://doi.org/10.1186/1479-5868-6-39>

- St-Amour, S., Cailhol, L., Ruocco, A. C., & Bernard, P. (2021). Could physical exercise be an effective treatment for adults with borderline personality disorder? *Psychiatry Research*, *295*, 113625. <https://doi.org/10.1016/j.psychres.2020.113625>
- St-Amour, S., Cailhol, L., Ruocco, A. C., & Bernard, P. (2022). Acute effect of physical exercise on negative affect in borderline personality disorder: A pilot study. *Clinical Psychology in Europe*, *4*(2). <https://doi.org/10.32872/cpe.7495>
- St-Amour, S., Bérubé, F., Cailhol, L., & Le Corff, C. (2022). Are physical activity and nutrition linked to personality disorders? Health habits and personality disorders: A scoping review. *Personality and Mental Health*, *17*(2), 147–156. <https://doi.org/10.1002/pmh.1568>
- Statistics Canada, Prince, S. A., Roberts, K. C., Reed, J. L., Biswas, A., Colley, R. C., & Thompson, W. (2020). *Daily physical activity and sedentary behaviour across occupational classifications in Canadian adults*. Government of Canada. <https://doi.org/10.25318/82-003-X202000900002-ENG>
- Stubbs, B., & Rosenbaum, S. (2018). In *Exercise-Based Interventions for Mental Illness: Physical Activity as Part of Clinical Treatment*. Academic Press.
- Vancampfort, D., De Hert, M., Stubbs, B., Soundy, A., De Herdt, A., Detraux, J., & Probst, M. (2015). A Systematic Review of Physical Activity Correlates in Alcohol Use Disorders. *Archives of Psychiatric Nursing*, *29*(4), 196–201. <https://doi.org/10.1016/j.apnu.2014.08.006>
- Vancampfort, D., Firth, J., Schuch, F. B., Rosenbaum, S., Mugisha, J., Hallgren, M., Probst, M., Ward, P. B., Gaughran, F., De Hert, M., Carvalho, A. F., & Stubbs, B. (2017). Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: a global systematic review and meta-analysis. *World Psychiatry*, *16*(3), 308–315. <https://doi.org/10.1002/wps.20458>
- Vancampfort, D., Hallgren, M., Firth, J., Rosenbaum, S., Schuch, F. B., Mugisha, J., Probst, M., Van Damme, T., Carvalho, A. F., & Stubbs, B. (2018). Physical activity and suicidal ideation: A systematic review and meta-analysis. *Journal of Affective Disorders*, *225*, 438–448. <https://doi.org/10.1016/j.jad.2017.08.070>
- Vancampfort, D., Knapen, J., Probst, M., Scheewe, T., Remans, S., & De Hert, M. (2011). A systematic review of correlates of physical activity in patients with schizophrenia. *Acta Psychiatrica Scandinavica*, *125*(5), 352–362. <https://doi.org/10.1111/j.1600-0447.2011.01814.x>
- Vancampfort, D., Probst, M., Scheewe, T., De Herdt, A., Sweers, K., Knapen, J., van Winkel, R., & De Hert, M. (2013). Relationships between physical fitness, physical activity, smoking and metabolic and mental health parameters in people with schizophrenia. *Psychiatry Research*, *207*(1-2), 25–32. <https://doi.org/10.1016/j.psychres.2012.09.026>
- Victor, S. E., & Klonsky, E. D. (2016). Validation of a Brief Version of the Difficulties in Emotion Regulation Scale (DERS-18) in Five Samples. *Journal of Psychopathology and Behavioral Assessment*, *38*(4), 582–589. <https://doi.org/10.1007/s10862-016-9547-9>
- WHO ASSIST Working Group. (2002). The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): development, reliability and feasibility. *Addiction*, *97*(9), 1183–1194. <https://doi.org/10.1046/j.1360-0443.2002.00185.x>
- Zhang, Z. (2016). Variable selection with stepwise and best subset approaches. *Annals of Translational Medicine*, *4*(7), 136–136. <https://doi.org/10.21037/atm.2016.03.35>